

Authors

Isabel Wang
BN (Hons), RN
School of Nursing and Midwifery, Griffith
University, Gold Coast, Queensland,
Australia

Dr Rachel Walker
PhD, RN
School of Nursing and Midwifery, Menzies
Health Institute, Griffith University,
Nathan Campus, Queensland, Australia
Division of Surgery, Princess Alexandra
Hospital, Queensland, Australia

Professor Brigid M Gillespie
PhD, RN, FACORN
School of Nursing and Midwifery, Griffith
University, Gold Coast, Queensland,
Australia
Gold Coast Hospital and Health Service,
Queensland, Australia

Corresponding author

Professor Brigid M Gillespie
PhD, RN, FACORN
School of Nursing and Midwifery, Griffith
University, Gold Coast, Queensland,
Australia
Gold Coast Hospital and Health Service,
Queensland, Australia
b.gillespie@griffith.edu.au

Pressure injury prevention for surgery: Results from a prospective, observational study in a tertiary hospital

Implementing pressure injury prevention in a perioperative setting

Background

Hospital acquired pressure injury (HAPI) refers to the development of pressure injury (PI) during hospitalisation. Prevalence rates of HAPI among all PI cases in acute health settings vary widely in different regions and countries, ranging from three to 18 per cent^{1–12}. While patients with limited mobility are at greater risk of developing a PI, anaesthetised patients are particularly vulnerable due to limited mobility. Perioperative HAPI remains problematic, with current prevalence rates varying from 5.1 to 64.1 per cent¹³. Through the literature review component of this study, we identified five categories of pressure injury prevention (PIP) strategies that comply with current clinical practice guidelines¹⁴. There has been considerable research undertaken on PIP in medical–surgical wards but research undertaken in the perioperative environment is scarce¹⁵.

Study aim

The aim of this study was to report on the PIP strategies used by perioperative health professionals at a large tertiary hospital in southeast Queensland.

Method

This is an observational study of the use of PIP strategies by health care professionals across a selected range of surgical procedures in a tertiary hospital. A structured data collection tool was developed, tested and used

to collect patients' demographic and clinical data, and health care professionals' implementation of PIP strategies, as well as data from skin inspection on day two after surgery. Observations occurred during the preoperative, intra-operative and post-operative periods, i.e. in the induction room, in the operating room and during the first half hour after the patient was admitted to the Post Anaesthesia Care Unit (PACU) respectively. The study was conducted in 2016.

Setting and sample

The study setting was a 750-bed tertiary hospital in Queensland. Surgical procedures were purposively selected across seven specialties. Adult patients whose length of surgery exceeded 60 minutes and who were anticipated to be an inpatient for a minimum of 48 hours after surgery were included in the study. All perioperative practitioners working in the operating room department were invited to participate. Patients who were unable to provide informed consent and who could not speak, read or understand English in the absence of an interpreter were excluded from the study.

Results

In total, this study recruited and observed 278 staff during surgical procedures of 73 recruited patients. Table 1 presents patients' characteristics by surgical specialty. Each specialty group had at least ten

Table 1: Patient characteristics by surgical specialty (n = 73)

Surgical specialty (Frequency and percentage of total patient sample)			Age (Median and IQR of total patient sample)		Gender (Frequency and percentage of specialty group)			Comorbidities (Median and IQR of total patient sample)		Waterlow risk groups [†] (Median and IQR of total patient sample)		Length of surgery (in minutes) (Median and IQR of total patient sample)		Temperature at PACU admission [^] (Median and IQR of total patient sample)	
	n	%	Median	IQR		n *	%	Median	IQR	Median ^{††}	IQR	Median	IQR	Median	IQR
Cardiac	10	13.7	60.0	21.75	male	8	80	4.5	7.25	4.00	1.25	302.5	141.75	36.40	0.00
Neurology	10	13.7	66.0	27.75	female	7	70	4.5	4.25	1.00	2.00	215.5	132.75	36.00	0.80
Orthopaedic	12	16.4	63.0	19.50	male	7	58	3.5	4.50	1.00	1.00	115.5	55.25	36.45	0.45
Gynaecology	11	15.1	52.0	22.00	female	11	100	4.0	1.75	1.00	4.00	181.0	62.00	36.10	0.40
General	10	13.7	75.5	18.50	male	6	60	4.0	4.25	1.00	3.00	268.5	167.00	36.30	1.05
Vascular	10	13.7	70.5	16.75	male	7	70	5.0	4.50	1.00	2.25	151.5	52.25	36.20	0.42
Urology	10	13.7	69.00	14.50	male	8	80	4.00	5.50	0.00	1.75	150.5	138.00	36.15	0.42

* n reports only the dominant gender per specialty group.

† 18 incomplete Waterlow risk assessments were excluded, as only 15 per cent of the items were complete, on average.

†† 1 to 3 refers to at-risk, moderate-risk and high-risk groups, respectively; 0 refers to not-at-risk group; 4 refers to unavailable or incomplete Waterlow assessment.

^ In 11 cases data for temperature upon PACU admission was missing, as this was not recorded when the patients arrived at the PACU.

patients, and their age, comorbidities, Waterlow risk assessment scores, length of surgery, and temperature upon PACU admission are reported using the median and IQR.

Data about selected PIP strategies were derived through observation or documentation; as shown in Table 2, most data was derived from direct observation. Data for three selected categories of PIP strategies – skin inspection, positioning aids and medical devices or/and equipment – were collected from direct observation, and data for selected interprofessional communication strategies were collected from both observation and documentation when communication was related to skin inspection, positioning and medical devices or/and equipment (n = 10/90, 55.6 per cent). Data for selected communication strategies where communication was related to

Table 2: Sources of data for each key PIP strategy category (n = 90)

Category of PIP strategies	Number of PIP strategies implemented*	Data for PIP strategy collected from observation n (%)	Data for PIP strategy collected from documentation n (%)
Skin inspection	6	6 (100)	0 (0)
Positioning aids	18	18 (100)	0 (0)
Medical devices or/and equipment	34	34 (100)	0 (0)
Thermal regulation	14	12 (85.7)	2 (14.3)
Interprofessional communication	18	8 (44.4)	10 (55.6)
Total number of PIP strategies	90	78	12

*Each category includes a number of PIP strategies. For instance, preoperative, intra-operative and post-operative skin integrity screening are three strategies in the skin inspection category.




Figure 1: Box plots of the total number of PIP strategies implemented during the intra-operative period with whiskers from minimum to maximum for the three Waterlow risk categories (n = 36)

thermal regulation was drawn from documentation when it was related to patients' temperature measured upon PACU admission (n = 2/90, 14.3%).

Figure 1 is a box plot that illustrates the number of PIP strategies implemented, based on patients' PI risk. Of the 36 patients deemed to be at risk of developing PI using the Waterlow risk assessments, the median number of PIP strategies implemented intra-operatively ranged from 39 to 40 strategies for the three risk groups – at risk, moderate risk, high risk (IQR = 11.5, 6.5 and 3, respectively). The results suggest that the highest number of PIP strategies was implemented for patients in the at-risk category. The minimum total number of PIP strategies implemented for patients in the moderate and high-risk groups was higher than for patients in the at-risk group, although the difference was not statistically significant (p = 0.819).

Sub-analysis

A high incidence rate of PI on day two after surgery has been reported in the literature^{16,17}. Hence, the decision to inspect patients' skin on the

second post-operative day was made. In this study, skin inspection on the second day after surgery revealed

that four male cardiac surgical patients had developed a post-operative Stage 1 PI on the ear, due to pressure from oxygen tubing.

As shown in Table 3 these patients had a median age of 70 years (IQR: 22 years, range: 55–79 years) and were all either overweight or obese. Two patients had more than eight comorbidities, and three had an American Society of Anesthesiologists (ASA) score of 4, indicating severe systemic disease that is a constant threat to life¹⁸. Prior to induction, a Waterlow risk assessment was not completed for three of these patients (75 per cent). The median length of surgery for this group was 302 minutes, over five hours (IQR = 162.5 minutes, range: 201–410 minutes). The primary surgical position used in all four cases was supine.

Table 3: Sub-analysis for patients who had PI on day two after surgery (n = 4)

Factors			n (%)
Demographic factors	Age	47–70 years	2 (50)
		> 71 years	2 (50)
	Number of comorbidities	0–3	1 (25)
		4–8	1 (25)
		> 8	2 (50)
	BMI	25.0–29.9	2 (50)
		≥ 30.0	2 (50)
	Waterlow scores	≥ 20	1 (25)
		N/A or incomplete	3 (75)
	ASA status	ASA 3	1 (25)
		ASA 4	3 (75)
Clinical factors	Number of medications taken before induction	0–2	3 (75)
		3–6	1 (25)
	Length of surgery	181–300 minutes	50
		> 300 minutes	50
	Primary surgical position *	supine	4 (100)

* The patient may be repositioned intra-operatively; however, the position in which the patient spent the most time was documented.

Key messages

The total number of PIP strategies used intra-operatively and Waterlow risk assessment scores suggests that patient deemed to be at risk (using the Waterlow risk assessment tool) have a higher number of PIP strategies implemented intra-operatively.

Patients having prolonged surgeries, i.e. greater than two hours, are at greater risk of developing a PI.

Post-operative PIs that occur as a result of using medical devices or equipment should not be overlooked. Perioperative professionals should be vigilant and monitor the use of devices and equipment during the intra-operative period.

References

1. Cuddigan J, DR Berlowitz, EA Ayello. (2001). Pressure ulcers in America: Prevalence, incidence, and implications for the future. *Adv Skin Wound Care* 2001;14(4):208–215.
2. Chan EY, Tan SL, Lee CK, Lee JY. Prevalence, incidence and predictors of pressure ulcers in a tertiary hospital in Singapore. *J Wound Care* 2005;14(8):383–384,386–388.
3. Reddy M, Gill SS, Rochon PA. Preventing pressure ulcers: A systematic review. *JAMA* 2006;296(8):974–984.
4. Victoria Department of Health. PUPPS3 – Pressure ulcer point prevalence survey: Statewide report 2006. Melbourne: Victoria Department of Health; 2006.
5. South Australia Department of Health. South Australia pressure ulcer point prevalence survey report 2007. Adelaide: South Australia Department of Health; 2007.
6. Sanada HY, Miyachi T, Ohura T, Moriguchi K, Tokunaga K, Shido et al. The Japanese pressure ulcer surveillance study: A retrospective cohort study to determine prevalence of pressure ulcers in Japanese hospitals. *Wounds* 2008;20(6):176–182.
7. VanGilder C, Amlung S, Harrison P, Meyer S. Results of the 2008–2009 international pressure ulcer prevalence survey and a three-year, acute care, unit-specific analysis. *Ostomy Wound Manage* 2009;55(11):39–45.
8. Mulligan S, Scott L, Prentice J, Elmes R, Carville K, Santamaria N. WoundsWest Wound prevalence survey 2009: State-wide report. Perth: Ambulatory Care Services, Department of Health; 2009.
9. Lyder CH, Wang Y, Metersky M, Curry M, Kliman R, Verzier NR. Hospital-acquired pressure ulcers: Results from the national Medicare Patient Safety Monitoring System study. *J Am Geriatr Soc* 2012;60(9):1603–1608.
10. Bergquist-Beringer SL, Dong L, He J, Dunton N. Pressure ulcers and prevention among acute care hospitals in the United States. *Jt Comm J Qual Patient Saf* 2013;39(9):404–414.
11. Mallah Z, Nassar N, Kurdahi Badr L. The effectiveness of a pressure ulcer intervention program on the prevalence of hospital acquired pressure ulcers: Controlled before and after study. *Appl Nurs Res* 2015;28(2):106–113.
12. Queensland Health. Queensland bedside audit. Brisbane: Queensland Health; 2016.
13. Shafipour V, Ramezanpour E, Gorji MA, Moosazadeh M. Prevalence of postoperative pressure ulcer: A systematic review and meta-analysis. *Electron Physician* 2016;8(11):3170–3176.
14. National Pressure Ulcer Advisory Panel (NPUAP), European Pressure Ulcer Advisory Panel (EPUAP) and Pan Pacific Pressure Injury Alliance (PPPIA). Prevention and treatment of pressure ulcers: Quick reference guide. Emily Haesler (Ed.). Cambridge Media: Perth, Australia; 2014.
15. Wang HM, Walker R, Gillespie BM. How well do perioperative practitioners implement pressure injury prevention guidelines? A brief report. *Wound Pract Res* 2018;26(1):24–33.
16. Schuurman JP, Schoonhoven L, Keller BP, van Ramshorst B. Do pressure ulcers influence length of hospital stay in surgical cardiothoracic patients? A prospective evaluation. *J Clin Nurs* 2009;18(17):2456–2463.
17. Connor T, Sledge JA, Bryant-Wiersema L, Stamm L, Potter P. Identification of preoperative and intra-operative variables predictive of pressure ulcer development in patients undergoing urologic surgical procedures. *Urol Nurs* 2010;30(5):289.
18. American Society of Anesthesiologists (ASA). ASA physical status classification system. Washington D.C.: ASA; 2014 [cited 2017 November 11]. Available from: www.asahq.org/resources/clinical-information/asa-physical-status-classification-system.